

Appln No. 10/510,617  
Amdt date January 26, 2009  
Reply to Office action of July 25, 2008

**Amendments to the Claims:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

**Listing of Claims:**

1-10 (Cancelled)

11. (Currently Amended) A pharmaceutical mixture for treating rheumatism, consisting of  
an alcoholic extract of:

*Tripterygium hypoglaucum* (Levl.) Hutch. 1-4 parts by weight;

*Epimedium brevicornum* Maxim. 1-4 parts by weight;

*Lycium barbarum* L. 1-4 parts by weight; and,

*Cuscuta chinensis* Lam., or *Cuscuta australis* R. Br., 1-4 parts by weight.

12. (Previously Presented) The pharmaceutical mixture according to claim 11, wherein

*Tripterygium hypoglaucum* (Levl.) Hutch. is 2 parts by weight;

*Epimedium brevicornum* Maxim. is 2 parts by weight;

*Lycium barbarum* L. is 1 part by weight; and,

*Cuscuta chinensis* Lam., or *Cuscuta australis* R. Br., 1 part by weight.

13. (Withdrawn) A method for preparing the pharmaceutical mixture according to claim 11,  
consisting essentially of:

providing and cutting the *Tripterygium hypoglaucum* (Levl.) Hutch into pieces;

providing and cutting the *Epimedium brevicornum* Maxim into pieces;

providing and optionally crushing the *Lycium barbarum* L;

Appln No. 10/510,617  
Amdt date January 26, 2009  
Reply to Office action of July 25, 2008

providing and optionally crushing the *Cuscuta chinensis* Lam;  
extracting the *Tripterygium hypoglaucum* (Levl.), *Epimedium brevicornum* Maxim,  
*Lycium barbarum* L, and *Cuscuta chinensis* Lam, separately or in combination, in a 0 to 95%  
alcohol/water mixture, at a temperature in the range of 1 to 98 °C, 1 to 4 times, to form one or  
more extracted liquor(s)/alcohol mixtures;  
optionally mixing the extracted liquor(s)/alcohol mixtures;  
recovering the extracted liquor(s) from the extracted liquor(s)/alcohol mixtures;  
condensing, drying, and smashing the extracted liquor(s) to form extracted herb powders;  
and,  
mixing the extracted herb powders uniformly or proportionally to form a pharmaceutical  
mixture.

14. (Withdrawn) A method for preparing the pharmaceutical mixture according to claim 11,  
consisting essentially of:

weighing the raw herbs according to the specified parts by weight;  
cutting *Epimedium brevicornum* Maxim into pieces;  
cutting the *Tripterygium hypoglaucum* (Levl.) Hutch into pieces;  
decocting the *Epimedium brevicornum* Maxim pieces with water three times;  
decocting the *Tripterygium hypoglaucum* (Levl.) Hutch pieces with water three times;  
immersing each of *Lycium barbarum* L or *Cuscuta chinensis* Lam in water at a  
temperature in the range of 80°C ~ 95°C for 1 to 3 times respectively;  
blending the decocted and immersed herbs to form a mixed fluid,  
loading the decocted or immersed fluid from each herb on a corresponding macroporous  
polymeric adsorbent column;

**Appln No. 10/510,617**  
**Amdt date January 26, 2009**  
**Reply to Office action of July 25, 2008**

absorbing each decocted or immersed fluid onto the corresponding macropore polymeric absorbent column;

washing each resin column with water until effluent becomes clear,

eluting each resin column with 30-99.5% alcohol to form an eluting liquor;

collecting the eluting liquor while the color ranges from a deep color to a very weak color;

forcing alcohol out of the column with water;

mixing the alcohol with the eluting liquor, wherein the weight of total eluent is 1 to 8 times the weight of the raw herbs;

recovering all 4 of the eluting liquors from the alcohol;

separately condensing all 4 of the eluting liquors to a specific gravity of 1.10;

spray drying each of the condensed eluting liquors to form extracted herb powders; and,

mixing the herb powders to form the pharmaceutical mixture.

15. (Previously Presented) The pharmaceutical mixture according to claim 11, wherein the mixture is in the form of a hard capsule, soft capsule, tablet, granule, or injectable liquid.

16. (Withdrawn) A method for preparing the pharmaceutical mixture according to claim 11, consisting essentially of:

providing and cutting *Tripterygium hypoglaucum* (Levl.) Hutch into pieces;

extracting from the *Tripterygium hypoglaucum* (Levl.) Hutch three times with water a decoction fluid, wherein the weight of the water used in the three extractions is, respectively, 13, 10, and 10 times the weight of the *Tripterygium hypoglaucum* (Levl.) Hutch prior to extraction, with each extraction lasting 1 hour;

**Appln No. 10/510,617**  
**Amdt date January 26, 2009**  
**Reply to Office action of July 25, 2008**

providing and cutting *Epimedium brevicornum* Maxim into segments;

extracting from the *Epimedium brevicornum* Maxim three times with water a decoction fluid, wherein the weight of the water used in the three extractions is, respectively, 15, 10, 10 times the weight of the *Epimedium brevicornum* prior to extraction, with each extraction lasting 1 hour;

crushing *Lycium barbarum* L to a powder and immersing the powder in 20 times by weight of water, at 80°C-95°C for 1 hour;

crushing *Cuscuta chinensis* Lam to a coarse powder and immersing the powder in 31 times by weight of water, at 90°C for 1 hour;

filtering the decoction fluid or immersion fluid of each of the *Tripterygium hypoglaucum* (Levl.), *Epimedium brevicornum* Maxim, *Lycium barbarum* L, and *Cuscuta chinensis* Lam separately;

pouring each filtered fluid through a macroporous polymeric adsorbent resin column;

eluting each column with 70% ethanol until the color of effluent becomes deep;

continuously collecting the effluent until the color of the effluent becomes very weak;

recovering the alcohol in the effluent of the *Tripterygium hypoglaucum* (Levl.),

*Epimedium brevicornum* Maxim, *Lycium barbarum* L, and *Cuscuta chinensis* Lam;

concentrating and drying the remaining fluid of the *Tripterygium hypoglaucum* (Levl.), *Epimedium brevicornum* Maxim, *Lycium barbarum* L, and *Cuscuta chinensis* Lam to form separate extract powders;

concentrating and drying the remaining fluid of the *Tripterygium hypoglaucum* (Levl.), *Epimedium brevicornum* Maxim, *Lycium barbarum* L, and *Cuscuta chinensis* Lam and,

mixing the extract powders uniformly and proportionally to form a pharmaceutical mixture.

**Appln No. 10/510,617**  
**Amdt date January 26, 2009**  
**Reply to Office action of July 25, 2008**

Claims 17-18 (Cancelled)